

[CLICK HERE TO JUMP TO USE DIRECTIONS](#)

[CLICK HERE TO JUMP TO BERMUDA & ZOYSIA DIRECTIONS](#)

[CLICK HERE TO JUMP TO BERMUDA & FESCUE DIRECTIONS](#)



# ORNAMEC®

OVER-THE-TOP\*

## GRASS HERBICIDE

### Controls Grass in Ornamentals, Nurseries & Landscaping.

#### ACTIVE INGREDIENT:

Fluazifop-P-butyl: Butyl(R)-2-[4-[[5-(trifluoromethyl)-2-

pyridinyl]oxy]phenoxy]propanoate ..... 6.75%

**INERT INGREDIENTS:** ..... 93.25%

TOTAL 100.00%

\*Contains 0.50 pound (+) isomer (fluazifop-P-butyl) per gallon.

Contains petroleum distillates, xylene or xylene range aromatic solvent.

ORNAMEC® is a registered trademark of PBI/GORDON CORPORATION.

### KEEP OUT OF REACH OF CHILDREN **CAUTION**

See below for additional Precautionary Statements and First Aid.



**READ THE ENTIRE LABEL FIRST.  
OBSERVE ALL PRECAUTIONS AND  
FOLLOW DIRECTIONS CAREFULLY.**

### PRECAUTIONARY STATEMENTS

#### Hazards to Humans & Domestic Animals

**CAUTION:** Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or on clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

### FIRST AID

If in eyes:	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
If on skin or clothing:	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
If swallowed:	<ul style="list-style-type: none"> <li>Immediately call a poison control center or doctor.</li> <li>Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>Do not give any liquid to the person.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>

(continued on next column)

### FIRST AID (continued)

#### If inhaled:

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.

**Note to Physician:** Contains petroleum distillate – vomiting may cause aspiration pneumonia.

#### Personal Protective Equipment (PPE):

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category E on an EPA chemical resistance category selection chart.

Applicators and other handlers (other than mixers and loaders) must wear: long-sleeved shirt and long pants, chemical resistant gloves such as barrier laminate, nitrile rubber, neoprene rubber, or Viton, and shoes plus socks.

Mixers and loaders must wear: long-sleeved shirt and long pants, chemical resistant gloves such as barrier laminate, nitrile rubber, neoprene rubber, or Viton, shoes plus socks, protective eyewear, and chemical resistant apron when mixing or loading.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### User Safety Recommendations:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

#### Environmental Hazards

This product is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater. Do not apply when weather conditions favor drift from target area.

#### Physical and Chemical Hazards

Do not use or store near heat or open flame.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of application.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR part 170.

This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

For early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear: coveralls, chemical resistant gloves such as barrier laminate, nitrile rubber, neoprene rubber, or Viton, and shoes plus socks.

## STORAGE & DISPOSAL

**PROHIBITIONS:** Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

**STORAGE:** Store in original container only. Keep container closed when not in use. Do not store near food or feed. In case of spill or leak on floor or paved surfaces, soak up with sand, earth or synthetic absorbent. Remove to chemical waste area.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

## GENERAL INFORMATION

ORNAMEC® is a selective postemergence herbicide for control of annual and perennial grass weeds. ORNAMEC does not control broadleaf weeds or sedges.

**Control Symptoms** — ORNAMEC is a systemic herbicide which moves from the treated foliage into the shoots, roots, rhizomes, stolons, and growing points (meristematic regions) of treated grass weeds. Excellent control of a wide range of annual and perennial grass weeds will be obtained with ORNAMEC when applied as recommended on this label.

Growth of treated grass weeds stops soon after application. Symptoms include loss of vigor, yellowing and/or reddening, and eventual death to the treated grass plant. Symptoms are generally observed within one to three weeks, depending on grass weed species and environmental conditions.

ORNAMEC can be used to control grass weeds in many newly transplanted and established ornamentals, trees, shrubs and ground-covers in noncrop areas. See the following Tables I through IV for specific recommended uses.

Application of ORNAMEC may be made over-the-top to ornamentals listed in Table I. Plants listed in Tables II through IV should have a directed spray applied to prevent contact of spray with foliage. If the growth habit of the plant or other factors do not allow a directed spray, an over-the-top application can be made to plants in Table II-IV. However, there is the possibility of damage in the range listed for each table when applied at label rates.

**Do not apply to ornamentals that may be harvested for food within one year after application.**

**Do not apply this product through any type or irrigation system.**

**NOTICE TO BUYER AND USER:** It is impossible to test every species and variety or cultivar of ornamental or nursery plants under all conditions. Plant tolerance of pesticides vary as conditions vary. Plant tolerance of ORNAMEC at label rates has been found to be acceptable within the ranges specified for the indicated genera and species. Neither the manufacturer nor the seller has determined whether or not ORNAMEC can be used on plants not specified on this label. The professional user should determine if ORNAMEC can be used prior to use.

## APPLICATION DIRECTIONS

**Timing** — For ornamental use apply to actively growing grass weeds before they exceed the recommended growth stages shown below.

Do not apply ORNAMEC herbicide if rainfall is expected within 1 hour.

**ALWAYS ADD A NONIONIC SURFACTANT** containing at least 75% surface wetting agent at 0.25% vol./vol. (½ pint per 25 gallons) of finished spray volume.

## SPOT TREATMENTS TO CONTROL GRASS IN ORNAMENTALS, NURSERIES AND LANDSCAPING AREAS

Mix ORNAMEC and only nonionic surfactant with water according to the amounts shown below. Spray to obtain thorough coverage, but do not spray to runoff. Re-treat if necessary.

To Make This Spray Volume	Add These Amounts*	
	ORNAMEC	Nonionic Surfactant
1 gal.	2.5 fl. oz. (4.5 tbsp.)	1/3 fl. oz.
10 gal.	26 fl. oz.	3 fl. oz.
25 gal.	64 fl. oz.	1/2 pint
50 gal.	128 fl. oz.	1 pint

\*1 fluid ounce (fl. oz.) = 2 tablespoons = 6 teaspoons

## ANNUAL AND PERENNIAL GRASS CONTROL RECOMMENDATIONS

- Use 64-96 fl. oz. of ORNAMEC herbicide per acre and 0.25% vol./vol. (½ pint per 25 gallons) of a nonionic surfactant. For perennial grasses, use 96 fl. oz. of ORNAMEC per acre per application and three applications will be needed.
- Spray to obtain thorough coverage of grass foliage, but not to runoff. Repeat treatments may be necessary to provide full season control.
- Use standard pesticide hollow cone or flat fan nozzles.
- Apply ORNAMEC herbicide to actively growing grasses before they exceed the recommended growth stages shown below.
- Do not apply ORNAMEC herbicide to grasses which are stressed due to moisture, temperature, low soil fertility, mechanical or chemical injury.
- Do not apply to grasses which have tillered, formed seedheads, or exceeded recommended growth stages.
- Apply ORNAMEC herbicide when the first grass species in a mixed grass weed population reaches the recommended growth stage for treatment.

## ADDITIONAL GRASS CONTROL

ORNAMEC may also be used to suppress and eventually control some undesirable grasses in turf areas.

To control common bermudagrass and hybrid bermudagrass, follow the directions below:

### BERMUDAGRASS CONTROL IN ZOYSIAGRASS AND TALL FESCUE TURF:

ORNAMEC may also be used to suppress and eventually control some undesirable grasses in turf areas. To control common bermudagrass and hybrid bermudagrass, follow the directions below:

#### Bermudagrass Control in Zoysiagrass:

Late Spring:	Apply 12-16 fluid ounces of this product per acre with a nonionic surfactant. This application should be made around June 1 and repeated at 28-30 days. For spot treatments use ⅓ fluid ounce (2 teaspoons) of this product and ⅓ fluid ounce (2 teaspoons) of nonionic surfactant in one gallon of water to treat 1,000 square feet.
Mid Summer:	During hot summer weather use 16-20 fluid ounces of product per acre with a nonionic surfactant. For spot treatments use 0.4 fluid ounces (2½ teaspoons) of this product and ⅓ fluid ounce (2 teaspoons) of nonionic surfactant in one gallon of water to treat 1,000 square feet.  <b>NOTE:</b> The 20 fluid ounce rate could cause temporary turf discoloration.
Late Summer:	When bermudagrass is preparing for dormancy use 8-12 fluid ounces of this product per acre with a nonionic surfactant. For spot treatments use 0.25 fluid ounces (1½ teaspoons) of this product with ⅓ fluid ounce (2 teaspoons) of nonionic surfactant in one gallon of water to treat 1,000 square feet.

#### Bermudagrass Control in Tall Fescue:

Apply 20-24 fluid ounces of product per acre. For spot treatment use 0.5 fluid ounce (1 tablespoon) of this product with ⅓ fluid ounce (2 teaspoons) of nonionic surfactant in one gallon of water to treat 1,000 square feet. The first application should be made in early spring (April/May) when bermudagrass is breaking dormancy. The second application should be during September/October when bermudagrass is



TABLE I (continued)

Common Name or Variety	Scientific Name
Croton	<i>Codiaeum variegatum</i>
Crown Vetch	<i>Vicia</i> sp.
Daylily	<i>Hemerocallis</i> hybrids
Debbie, Jennie, Mahogany, Stargaze	<i>Gladiolus x hortulanus</i>
Delaware Valley white	<i>Rhododendron</i> sp.
Densata	<i>Picea glauca</i>
Dense yew	<i>Taxus x media</i>
Desert fern tree	<i>Lysiloma thornberii</i>
Desert gum	<i>Eucalyptus rudis</i>
Desert willow	<i>Pittosporum phillyraeoides</i>
Doublefile viburnum/tomentosum	<i>Viburnum plicatum</i>
Douglas fir	<i>Pseudotsuga menziesii</i>
Dusty miller	<i>Centaurea cineraria</i>
Dwarf Alberta	<i>Picea glauca</i>
Dwarf burford holly	<i>Ilex cornuta</i>
Dwarf Ninebark, Nanus	<i>Physocarpus opulifolius</i>
Dwarf Swiss mountain pine	<i>Pinus mugo</i>
Dwarf yaupon/Tall yaupon	<i>Ilex vomitoria</i>
Eastern hemlock	<i>Tsuga canadensis</i>
Eastern white pine	<i>Pinus strobus</i>
Elizabeth Gable	<i>Rhododendron "Gable Hybrid"</i>
Ellen Danica, grape ivy	<i>Cissus rhombifolia</i>
Emerald green	<i>Thuja occidentalis</i>
Emerald mound	<i>Lonicera xylosteum</i>
English ivy	<i>Hedera helix</i>
English roseum	<i>Rhododendron catawbiense</i>
Escallonia	<i>Escallonia fradessii</i>
Ethiopia banana	<i>Escallonia rubra</i>
Euonymus	<i>Musa maurelli</i>
Exotica weeping fig	<i>Euonymus fortunei</i>
False spirea	<i>Ficus benjamina</i>
Fashion	<i>Astilbe x arendsi</i>
Fastadiata	<i>Rhododendron</i> sp.
Fatshedera	<i>Erythrina fusca</i>
Feathery cassia	<i>Fatschedera lizei</i>
Fernleaf yarrow	<i>Cassia artemisioides</i>
Firecracker	<i>Achillea filipendulina</i>
Firewheel tree	<i>Justicia spicigera</i>
Flaviramea	<i>Stenocarpus sinuatus</i>
Flossflower, Pussyfoot	<i>Cornus sericea</i>
Flowering almond	<i>Ageratum</i> sp.
Flowering dogwood	<i>Prunus triloba</i>
Forsythia	<i>Cornus florida</i>
Forsythia	<i>Forsythia intermedia</i>
Forsythia	<i>Forsythia</i> sp.
Fosteri holly	<i>Ilex x attenuata</i>
Fransher evergreen	<i>Aglaonema commutatum</i>
Froebeilia	<i>Spiraea x bumalda</i>
Gardenia, dwarf	<i>Gardenia jasminoides</i>
Gay feather	<i>Liatris spicata</i>
Gazania	<i>Gazania uniflora leucolaena</i>
Gazania gold rush	<i>Gazania splendens</i>
Geranium	<i>Pelargonium domesticum</i>
Gerard's rose	<i>Rhododendron</i> sp.
Giant bird of paradise	<i>Strelitzia nicolai</i>
Giant dumbcane	<i>Dieffenbachia amoena</i>
Gibraltar	<i>Rhododendron</i> sp.
Globosa	<i>Thuja occidentalis</i>
Gloria	<i>Rhododendron</i> sp.
Glossy abelia	<i>Abelia grandiflora</i>
Golden bamboo	<i>Phyllostachys aurea</i>
Golden fruited palm	<i>Chrysalidocarpus lutescens</i> (small)
Gold coast juniper	<i>Juniperus chinensis</i>
Gold drop	<i>Physocarpus fruiticosus</i>
Gold drop, Primrose beauty	<i>Potentilla fructosa</i>
Gold flame	<i>Spiraea x bumalda</i>
Gold tip	<i>Juniperus chinensis</i>
Grapefruit	<i>Citrus paradisi</i>
Grayleafed Euryops	<i>Euryops pectinatus</i>
Green liriope/Variegated liriope	<i>Liriope muscari</i>
Green palo verde	<i>Parkinsonia aculeata</i>
Green stone crop	<i>Sedum brevifolium</i>
Greeting	<i>Rhododendron</i> sp.
Gumpo pink	<i>Rhododendron</i> sp.
Gumpo white	<i>Rhododendron</i> sp.

TABLE I (continued)

Common Name or Variety	Scientific Name
Gum-barked coolibah	<i>Eucalyptus microtheca</i>
Hahnii/Mother-in-law's tongue	<i>Sansevieria trifasciata</i>
Hahn red	<i>Rhododendron</i> sp.
Hahn's ivy	<i>Hedera helix hahnii</i>
Hedgehog cactus	<i>Echinocactus</i> sp.
Hen and chickens	<i>Sempervivum tectorum</i>
Herbert	<i>Rhododendron</i> sp.
Hicks yew	<i>Taxus x media</i>
Hino red	<i>Rhododendron</i> sp.
Hinoki falsecypress	<i>Chamaecyparis obtusa</i>
Hollyhock	<i>Alcea rosea</i>
Hollywood	<i>Juniperus chinensis</i>
Honey locust/shade master	<i>Gleditsia triacanthos</i> var. <i>inermis</i>
Hybrid tea rose	<i>Rosa hybrida</i>
H. H. Hume	<i>Rhododendron</i> sp.
Indian laurel	<i>Ficus macrocarpa nitida</i>
Indian laurel	<i>Iris</i> sp.
Iris	<i>Olenea tesota</i>
Ironwood	<i>Cupressus sempervirens</i>
Italian cypress	<i>Pelargonium peltatum</i>
Ivy geranium	<i>Ixora coccinea</i>
Ixora	<i>Jacaranda acutifolia</i>
Jacaranda	<i>Physocarpus fruticosus</i>
Jackmanni	<i>Syringa villosa</i>
James McFarlane lilac	<i>Buxus microphylla</i> var. <i>japonica</i>
Japanese boxwood	<i>Ilex crenata</i>
Japanese holly	<i>Pachysandra terminalis</i>
Japanese pachysandra	<i>Taxus cuspidata</i>
Japanese yew	<i>Aucuba japonica variegata</i>
Japanese laurel	<i>Acacia schafnerii</i>
Jim wheat acacia	<i>Simmondsia chinensis</i>
Joboba	<i>Viburnum x juddii</i>
Judd	<i>Juniperus chinensis</i>
Juniper	<i>Juniperus procumbens</i>
Juniper	<i>Juniperus sabina</i>
Kaempo	<i>Rhododendron</i> sp.
Kaffir lily	<i>Clivia miniata</i>
Keteleeri	<i>Juniperus chinensis</i>
Kluis sensation	<i>Rhododendron</i> sp.
Korean azalea/Poukhanense	<i>Rhododendron yedoense</i>
Korean boxwood	<i>Buxus microphylla</i> var. <i>koreana</i>
Korean lilac	<i>Syringa patula</i>
Lavender cotton	<i>Santolina chamaecyparisss</i>
Lawson cypress	<i>Chamaecyparis lawsoniana</i>
Lees dark purple	<i>Rhododendron catawbiense</i>
Lily-of-the-valley bush	<i>Pieris japonica</i>
Liriopae	<i>Liriope spicata</i>
Lodense pyracantha	<i>Pyracantha koidzumii</i>
Madwort	<i>Alyssum</i> sp.
Mahonia	<i>Mahonia aquifolium</i>
Manila Ripple	<i>Schefflera arboricola</i>
Maple/japanese	<i>Acer palmatum</i>
Marigold	<i>Tagetes</i> sp.
Marigold	<i>Calendula</i> sp.
Masasoit	<i>Rhododendron</i> sp.
Mediterranean fan palm	<i>Chamaerops humilis</i>
Mentor barberry	<i>Berberis mentorensis</i>
Meserve Holly	<i>Ilex meserveae</i>
Mexican border pine	<i>Pinus strobiformis</i>
Mexican fan palm	<i>Washingtonia robusta</i>
Mexican indigo	<i>Justicia spicigera</i>
Micans velvetleaf philodendron	<i>Philodendron oxycardium</i>
Mirror plant	<i>Coprosma baueri</i>
Mojave firethorn	<i>Pyracantha koidzumii</i> x <i>coccinea</i>
Moneymort/Wandering Sally	<i>Lysimachia nummularia</i>
Moon Glow	<i>Sansevieria</i> sp.
Morrow honeysuckle	<i>Lonicera x morrowii</i>
Mother's Day	<i>Rhododendron</i> sp.
Myres asparagus	<i>Asparagus densiflorus</i>
Myrtle, Dwarf periwinkle	<i>Vinca minor</i>
Nanum	<i>Viburnum opulus</i>
Narrow-leaf gimlet	<i>Eucalyptus spathulata</i>
Natal plum	<i>Carissa grandiflora</i>

TABLE I (continued)

Common Name or Variety	Scientific Name
Newport red	<i>Weigela florida</i>
New Zealand christmas tree	<i>Metrosideros excelsus</i>
Nicks compact juniper	<i>Juniperus chinensis</i>
Noble fir	<i>Abies procera</i>
Norfolk Island pine	<i>Araucaria heterophylla</i>
Norway maple	<i>Acer platanoides</i>
Norway spruce	<i>Picea abies</i>
Oakleaf hydrangea	<i>Hydrangea quercifolia</i>
Oak, live	<i>Quercus virginiana</i>
Ocotillo	<i>Fouquieria splendens</i>
Olive tree	<i>Olea europaea</i>
Oregon Grapeholly	<i>Magnolia sp.</i>
Ornamental strawberry	<i>Fragaria chiloensis</i>
Osmanthus, tea olive	<i>Osmanthus fragrans</i>
O'Connors legume	<i>Trifolium fragiferum</i>
Pagoda flower	<i>Clerodendrum speciosum</i>
Painted evergreen	<i>Aglanoema crispum</i>
Palibin	<i>Syringa meyeri</i>
Panicle hydrangea	<i>Hydrangea paniculata</i>
Parsley panax	<i>Polyscias fruticosa</i>
Parsoni	<i>Juniperus chinensis</i>
Passion vine	<i>Passiflora pfordtii</i>
Pepin	<i>Juniperus sabina</i>
Pericat	<i>Rhododendron sp.</i>
Periwinkle	<i>Vinca major</i>
Peter pan lily of the Nile	<i>Agapanthus africanus</i>
Petunia	<i>Petunia sp.</i>
Philodendron	<i>Philodendron selloum</i>
Photinia	<i>Photinia x fraseri</i>
Pink clover	<i>Polygonum capitatum</i>
Pink lady	<i>Raphiolepis indica</i>
Pink pearl	<i>Rhododendron sp.</i>
Pink weigelia	<i>Weigela florida</i>
Pink, variegated, petite oleander	<i>Nerium oleander</i>
Plumosa	<i>Juniperus horizontalis</i>
President Lincoln	<i>Rhododendron sp.</i>
Privet/california	<i>Ligustrum ovalifolium</i>
Prize	<i>Rhododendron sp.</i>
Prostrate myoporum	<i>Myoporum parvifolium</i>
Purple elegans	<i>Rhododendron catawbiense</i>
Purple false eranthemum	<i>Pseuderanthemum atropurpureum</i>
Purple Gem	<i>Rhododendron sp.</i>
Purple hophseed bush	<i>Dodonea viscosa purpurea</i>
Purple lantana (trailing)	<i>Lantana sellowiana</i>
Purple splendor	<i>Rhododendron</i>
Purple trailing ice plant	<i>Mesembryanthemum lampranthus</i>
Pygmy date palm	<i>Phoenix roebelenii</i>
Pyracantha	<i>Pyracantha gruberi</i>
Pyramidalis	<i>Thuja occidentalis</i>
Queen palm	<i>Acrecastrum romanoffianum</i>
Queen palm	<i>Cocos plumosa</i>
Radiator plant	<i>Peperomia scandens</i>
Raspberry ice	<i>Bougainvillea sp.</i>
Red bearberry	<i>Arctostaphylos uva-ursi</i>
Red box gum	<i>Eucalyptus polyanthemos</i>
Red bud	<i>Cercis canadensis</i>
Red cedar	<i>Juniperus virginiana</i>
Red fountain grass	<i>Pennisetum setaceum</i>
Red gum	<i>Eucalyptus rostrata</i>
Red pine	<i>Pinus resinosa</i>
Red ruffle	<i>Rhododendron sp.</i>
Red spike ice plant	<i>Mesembryanthemum spectabilis</i>
Red twig dogwood	<i>Cornus sericea</i>
Red wing	<i>Rhododendron sp.</i>
Rhododendron	<i>Rhododendron formosa</i>
Road runner	<i>Rhododendron sp.</i>
Rock rose	<i>Cistus hybridus</i>
Rose	<i>Rosa sp.</i>
Rosebud	<i>Rhododendron sp.</i>
Rosemary dwarf	<i>Rosmarinus officinalis prostratus</i>
Roseum elegans	<i>Rhododendron catawbiense</i>
Roseum superbum	<i>Rhododendron catawbiense</i>
Rose greely	<i>Rhododendron</i>

TABLE I (continued)

Common Name or Variety	Scientific Name
Rose ice plant	<i>Mesembryanthemum drosanthemum hispidum</i>
Royalty	<i>Rhododendron sp.</i>
Royal beauty	<i>Cotoneaster dammeri</i>
Rubber tree	<i>Ficus elastica decora</i>
Russian olive	<i>Elaeagnus angustifolia</i>
Rutherfordiana Constance	<i>Rhododendron sp.</i>
Sago palm	<i>Cycas revoluta</i>
Saguaro cactus	<i>Carnegiea gigantea</i>
Salmon spray	<i>Rhododendron sp.</i>
Saltbrush	<i>Atriplex sp.</i>
Sandankwa viburnum	<i>Viburnum suspensum</i>
Sandwort	<i>Arenaria verna</i>
Sargent juniper	<i>Juniperus chinensis</i>
Sasanqua camellia	<i>Camellia sasanqua</i>
Sawara cypress	<i>Chamaecyparis pisifera</i>
Scarlet firethorn, Lalandei	<i>Pyracantha coccinea</i>
Scotch heather	<i>Calluna vulgaris</i>
Scotch pine	<i>Pinus sylvestris</i>
Sedum	<i>Sedum spectabile</i>
Sedum	<i>Sedum x rubrotinctum</i>
Senna	<i>Cassia sturtii</i>
Serbian spruce	<i>Picea omorika</i>
Shasta daisy	<i>Chrysanthemum x superbum</i>
Shield fern	<i>Polystichum capense</i>
Shoe-string acacia	<i>Acacia stenophylla</i>
Shore juniper	<i>Juniperus conferta</i>
Showy crabapple	<i>Malus floribunda</i>
Siebold	<i>Euonymus alata</i>
Silk oak	<i>Grevillea robusta</i>
Silver King	<i>Euonymus japonica</i>
Silver maple	<i>Acer saccharinum</i>
Silver Queen evergreen	<i>Aglanoema commutatum</i>
Skandia	<i>Juniperus sabina</i>
Slender deutzia	<i>Deutzia gracilis</i>
Snow	<i>Rhododendron sp.</i>
Snowmound spirea	<i>Spiraea nipponica</i>
Snow in summer	<i>Cerastium tomentosum</i>
Sour orange	<i>Citrus aurantium</i>
Spanish dagger	<i>Yucca gloriosa</i>
Spider flower	<i>Grevillea rosmarinifolia</i>
Spreading cotoneaster	<i>Cotoneaster divaricatus</i>
Spreading Euonymus	<i>Euonymus kiautschowica</i>
Sprengeri	<i>Asparagus densiflorus</i>
Sprenger asparagus	<i>Asparagus densiflorus</i>
Star jasmine	<i>Trachelospermum jasminoides</i>
Star Magnolia	<i>Magnolia stellata</i>
Stewartstonian	<i>Rhododendron sp.</i>
Swamp immortella	<i>Erythrina fusca</i>
Sweetheart	<i>Rhododendron sp.</i>
Sweet William	<i>Dianthus barbatus</i>
Sword fern	<i>Nephrolepsis exaltata</i>
Tabor	<i>Rhododendron sp.</i>
Tahitian gardenia	<i>Gardenia taitensis</i>
Tall hedge buckthorn	<i>Rhamnus frangula</i>
Techny	<i>Thuja occidentalis</i>
Techny american arborvitae	<i>Thuja occidentalis</i>
Texas privet	<i>Ligustrum texanum</i>
Texas sage	<i>Leucophyllum frutescens</i>
Thayeri yew	<i>Taxus x media</i>
Threadleaf coreopsis	<i>Coreopsis verticillata</i>
Ti	<i>Cordyline terminalis</i>
Tortuosa corkscrew willow	<i>Salix matsudana</i>
Torulosa	<i>Juniperus chinensis</i>
Tradition	<i>Rhododendron sp.</i>
Trebubii Ribbon evergreen	<i>Aglanoema commutatum</i>
Variegated firethorn	<i>Pyracantha angustifolia</i>
Variegated hosta	<i>Hosta lancifolia</i>
Variegated mirror plant	<i>Coprosma repens</i>
Viburnum	<i>Viburnum odoratissimum</i>
Viburnum/japanese snowball	<i>Viburnum dentatum</i>
Viburnum/willowwood	<i>Viburnum x rhytidophyloides</i>
Vicari	<i>Ligustrum x Vicari</i>
Waffle plant/Metallic plant	<i>Hemigraphis sp.</i>
Water willow	<i>Jacobinia ghiesbrechtiana</i>
Wax myrtle	<i>Myrica cerifera</i>
Weeping dagger	<i>Yucca pendula</i>

**TABLE I (continued)**

Common Name or Variety	Scientific Name
Weeping forsythia	<i>Forsythia suspensa</i>
Weeping willow	<i>Salix alba</i>
Welleri	<i>Buxus sempervirens</i>
Western/Ponderosa pine	<i>Pinus ponderosa</i>
Wheelers dwarf, variegated	<i>Pittosporum tobira</i>
White african daisy	<i>Osteospermum fruticosum alba</i>
White cascade	<i>Rhododendron sp.</i>
White catalpa	<i>Rhododendron catawbiense</i>
White cedar	<i>Thuja occidentalis</i>
Willowleaf cotoneaster	<i>Cotoneaster salicifolius franch</i>
Willow acacia	<i>Acacia saligna</i>
Willow-leaved acacia	<i>Acacia salacina</i>
Windmill palm	<i>Chamaerops excelsa</i>
Wichita blue	<i>Juniperus scopulorum</i>
Woodwardii	<i>Thuja occidentalis</i>
Xylosma	<i>Xylosma senticosa</i>
Yedda/Indian hawthorn	<i>Raphiolepis umbellata</i>
Yellow bells	<i>Tecoma stans angustate</i>
Yellow oleander tree	<i>Thevetia peruviana</i>
Yew pine	<i>Podocarpus macrophylla</i>
Yucca	<i>Yucca filamentosa</i>
Zinnia sp.	

**TABLE II**

DIRECTED APPLICATIONS are recommended to minimize the foliar injury of the ornamentals shown in the list below:

- When possible and plant growth habit allows, applications should be made as a **directed spray** to the plants listed below.
- Limited testing of the same plants has shown phytotoxicity of up to 20% when ORNAMEC herbicide is applied **over-the-top** at label rates.

Common Name or Variety	Scientific Name
Arcadia	<i>Gamolepsis chrysanthemoides</i>
Blue Pacific	<i>Juniperus sabina</i>
Blue rug	<i>Juniperus conferta</i>
Broadmoor	<i>Juniperus horizontalis</i>
Bugle weed	<i>Juniperus sabina</i>
Cats claw, Yellow trumpet	<i>Ajuga variegata</i>
Chinese snowball	<i>Begonia tweediana</i>
Columbine	<i>Viburnum macrocephalum</i>
Eureka	<i>Aquilegia hybrida</i>
Fetterbush	<i>Rhododendron obtusum</i>
Fraser Fir	<i>Leucothoe axillaris</i>
Gallery	<i>Abies fraseri</i>
Gazania	<i>Gladiolus x hortulanus</i>
Green carpet	<i>Gazania rigens</i>
Green fountain grass	<i>Herniaria glabra</i>
Grey owl	<i>Pennisetum setaceum</i>
Heavenly bamboo	<i>Juniperus virginiana</i>
Hughes	<i>Nandina domestica</i>
Indica	<i>Juniperus horizontalis</i>
Japanese honeysuckle	<i>Rhododendron indicum</i>
Japanese privet	<i>Lonicera japonica</i>
Juniper	<i>Ligustrum japonicum</i>
Kurume	<i>Juniperus japonica</i>
Lavender star plant	<i>Rhododendron obtusum</i>
Lemon-scented gum	<i>Grewia caffra</i>
Leyland cypress	<i>Eucalyptus citriodora</i>
Lilac	<i>Cupressocyparis leylandii</i>
Maki	<i>Syringa chinensis</i>
Maney	<i>Podocarpus macrophyllus</i>
Massangeana, dracaena	<i>Juniperus chinensis</i>
Mondo grass	<i>Dracaena fragans</i>
Nana	<i>Ophiopogon japonicum</i>
Old Gold	<i>Juniperus chinensis</i>
Oleander	<i>Juniperus chinensis</i>
Oyster plant	<i>Nerium oleander standard</i>
Pathfinder	<i>Rhoeo spathacea</i>
Pfitzeriana	<i>Juniperus scopulorum</i>
Philodendron	<i>Juniperus chinensis</i>
Pineapple guava	<i>Philodendron sp.</i>
Prickly pear cactus	<i>Feijoa sellowiana</i>
Prostrata	<i>Opuntia sp.</i>
P.J.M.	<i>Juniperus chinensis</i>
Red Maple	<i>Rhododendron sp.</i>
Redroot	<i>Acer rubrum</i>

**TABLE II (continued)**

Common Name or Variety	Scientific Name
Robusta	<i>Juniperus chinensis</i>
Ruellia	<i>Ruellia ciliosa</i>
Scandia	<i>Juniperus sabina</i>
Sky Rocket	<i>Juniperus virginiana</i>
Spearmint	<i>Juniperus chinensis</i>
Spring cinquefoil	<i>Potentilla verna</i>
Strawberry tree	<i>Arbutus unedo</i>
Tamariseifolia	<i>Juniperus sabina</i>
Temple tree	<i>Plumeria acuminata</i>
Tricolor dracaena	<i>Dracaena marginata</i>
Vanhoutte spirea	<i>Spiraea x vanhouttei</i>
Variagata	<i>Juniperus horizontalis</i>
Variegated ajuga	<i>Ajuga reptans</i>
Webberi	<i>Juniperus horizontalis</i>
Weeping bottle-brush	<i>Callistemon viminalis</i>
Welchii	<i>Juniperus scopulorum</i>
White lantana	<i>Lantana montevidensis</i>
Willow	<i>Salix caroliniana</i>
Wiltonii	<i>Juniperus horizontalis</i>
Youngtown Compacta	<i>Juniperus horizontalis</i>

**TABLE III**

DIRECTED APPLICATIONS are recommended to minimize the foliar injury of the ornamentals shown in the list below:

- When possible and plant growth habit allows, applications should be made as a **directed spray** to the plants listed below.
- Limited testing of the same plants has shown phytotoxicity of up to 50% when ORNAMEC herbicide is applied **over-the-top** at label rates.

Common Name or Variety	Scientific Name
Acacia	<i>Acacia latifolia</i>
Armstrongii	<i>Juniperus chinensis</i>
Baby rubber plant	<i>Peperomia obtusifolia</i>
Bleeding heart	<i>Dicentra spectabilis</i>
Bottle tree	<i>Brachychiton populneus</i>
Burkii	<i>Juniperus virginiana</i>
Carrot wood	<i>Cupaniopsis anacardioipes</i>
Chinese crenate croton	<i>Exococaria cochinchinensis</i>
Cordyline	<i>Cordyline stricta</i>
Coromandel	<i>Asystasia gangetica</i>
Desert broom	<i>Baccharis sarothroides</i>
Eucalyptus	<i>Eucalyptus nicholii</i>
Excelsa Striata	<i>Juniperus scopulorum</i>
Fiddlewood	<i>Citharexylum spinosum</i>
Formosa	<i>Rhododendron indicum</i>
Fragrant sumac	<i>Rhus aromatica</i>
Hearts and flowers	<i>Aptenia cordifolia</i>
Hersey red	<i>Rhododendron obtusum</i>
Hibiscus	<i>Hibiscus lepenk</i>
Hinodegeri	<i>Rhododendron obtusum</i>
Hino pink	<i>Rhododendron obtusum</i>
Jade Plant	<i>Crassula argentea</i>
Janet Craig/Warneckii	<i>Dracaena deremensis</i>
Karen	<i>Rhododendron poukhanense</i>
Kings crown	<i>Justicia carnea</i>
Laurel-leaved thunbergia	<i>Thunbergia laurifolia</i>
Marble Queen pothos	<i>Epipremnum aureum</i>
Mexican evening primrose	<i>Oenothera berlandieri</i>
Pinkhead knotweed	<i>Polygonum capitatum</i>
Red justicia	<i>Odontonema strictum</i>
Senna	<i>Cassia condolioma</i>
Shrimp plant	<i>Justicia brandegeana</i>
Slipper flower	<i>Pedilanthus tithymaloides</i>
Sonoran palo verde	<i>Cercidium praecox</i>
Southern magnolia	<i>Magnolia grandiflora</i>
Spiny Greek	<i>Juniperus scopulorum</i>
Swedish ivy	<i>Plectranthus australis</i>
Sweet acacia	<i>Acacia farnesiana</i>
Tifblue Blueberry	<i>Vaccinium Ashei</i>
Umbrella plant	<i>Cyperus alternifolius</i>
White ice plant (trailing)	<i>Mesembryanthemum delosperma alba</i>
White shrimp plant	<i>Justicia betonica</i>
Yellow shrimp plant	<i>Pachystachys lutea</i>

**TABLE IV**

DIRECTED APPLICATIONS are recommended to minimize the foliar injury of the ornamentals shown in the list below:

- When possible and plant growth habit allows, applications should be made as a **directed spray** to the plants listed below.
- Limited testing of the same plants has shown phytotoxicity greater than 50% when ORNAMEC herbicide is applied **over-the-top** at label rates.

Common Name or Variety	Scientific Name
Australia tree fern	Alsophila australis
Bar Harbor	Juniperus horizontalis
Birch/River	Betula nigra
Blue Chip	Juniperus horizontalis
Blue Haven	Juniperus scopulorum
Boulevard falsecypress	Chamaecyparis pisifera
Chandelier plant	Kalanchoe tubiflora
Compacta	Euonymus alata
Copperleaf	Acalypha godseffiana hertophylla
Hinocrimson	Rhododendron obtusum
Katherine Dykes	Physocarpus fruticosa
Lavender-scallops	Kalanchoe fedtschenkoi
Madagascar periwinkle	Catharanthus roseus
Pampas grass	Cortaderia selloana
Prince of Wales	Juniperus horizontalis
Purple heart	Setcreasea purpurea
Sea Green	Juniperus chinensis
Spider plant	Chlorophytum comosum
Wandering jew	Zebrina pendula

### LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

**THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED.** This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

778/9-2004 AP052504  
EPA REG. NO. 2217-728

Manufactured By  
 **pbi/gordon**  
**corporation**  
*An Employee-Owned Company*  
**1217 West 12th Street**  
**Kansas City, Missouri 64101**  
[www.pbigordon.com](http://www.pbigordon.com)