

Drug Facts (continued)

Inactive ingredients (continued)

D&C Yellow No. 10, FD&C Blue No. 1, gelatin, glycerin, methylparaben, polyethylene glycol, potassium propylene glycol, propylparaben, purified water, sorbitol special, sorbitol solution, xanthan gum

Questions or comments? Call toll-free 1-844-832-1138 Monday through Friday 9AM - 5PM EST or www.healthlifeusa.com

REMAIN CARTON FOR COMPLETE DRUG FACTS

*This product is not recommended or distributed by Procter & Gamble, the distributor of NyQuil® LiquiCaps®



NDC 69517-103-12

Compare to NyQuil® LiquiCaps® active ingredients*

Multi-Symptom Nighttime Cold & Flu

Acetaminophen 325 mg / Pain Reliever / Fever Reducer
Dextromethorphan HBr 15 mg / Cough Suppressant
Doxylamine Succinate 6.25 mg / Antihistamine

12 SOFTGELS

SEE NEW WARNINGS INFORMATION

Distributed by: HealthLife of USA LLC
Rahway, NJ 07065
www.healthlifeusa.com



0 69517 10312 3

Lot. No.:
Exp. Date:

Drug Facts (continued)

■ new symptoms occur
■ redness or swelling is present
■ cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults & for children even if you do not notice any signs or symptoms.

Directions

■ take only as directed - see Liver warning
■ do not exceed 8 doses per 24 hours
■ adults and children 12 years of age and over: 2 softgels with water every 4 hrs
■ children 4 to under 12 years of age: ask a doctor
■ children under 4 years of age: do not use
■ when using other Daytime or Nighttime products, carefully read each label to insure correct dosing

Other information

■ store at controlled room temperature 20°-25°C (68°-77°F)
■ read all product information before using
■ TAMPER EVIDENT DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts (continued)

prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or for emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

■ do not take if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

■ liver disease ■ glaucoma ■ diabetes
■ a breathing problem such as emphysema or chronic bronchitis ■ persistent or chronic cough as occurs with smoking, asthma, or emphysema
■ trouble urinating due to an enlarged prostate gland
■ cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

■ taking sedative or tranquilizers
■ taking the blood thinning drug warfarin

When using this product

■ excitability may occur, especially in children
■ be careful when driving a motor vehicle or operating machinery ■ avoid alcoholic drinks
■ alcohol, sedatives, and tranquilizers may increase drowsiness
■ may cause marked drowsiness

Stop use and ask a doctor

■ you get nervous, dizzy or sleepless
■ pain or cough gets worse or lasts more than 7 days
■ fever gets worse or lasts more than 3 days

Drug Facts (continued)

Active ingredients Purpose (in each softgel)

Acetaminophen 325 mg... Pain reliever/fever reducer
Dextromethorphan HBr 15 mg... Cough suppressant
Doxylamine succinate 6.25 mg... Antihistamine

Uses

■ temporarily relieves common cold/flu symptoms:
■ runny nose and sneezing ■ sore throat
■ fever ■ cough due to minor throat and bronchial irritation ■ headache ■ minor aches and pains


Warnings


Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 8 softgels (2,600 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

Sore throat warning: If sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use: ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a


DESIGN DEPARTMENT, INC. - 716-873-5008


 DIE LINE DO NOT PRINT

 VARNISH AREA

Date To Disk: 5/20/16


Job No.: 9462

 CMYK

 pms 293 Pantone Process Blue

Job prints 4 color process, please match

If Design Department, Inc. does not have an opportunity to view a "HARD COPY" or "PAPER" proof we will not be responsible for incorrect colors or printing errors in final printing. WE DO NOT CONSIDER ANY ELECTRONIC FILES INCLUDING PDF FILES AS A HARD COPY OR PAPER PROOF.

 **DRUG FACTS FORMAT WARNING**

Title - 9pt
Headings - 8pt
Body Copy - 5.8pt

We experienced problems formatting the Drug Facts information into the allotted space. Please note the above concessions to the minimum allowable text sizes.