



HealthLife
Naproxen Sodium Caplets
 Pain Reliever / Fever Reducer (NSAID)
 Strength to Last 12 Hours
 RETAIN CARTON FOR COMPLETE DRUG FACTS
 †This product is not manufactured or distributed by Bayer Healthcare LLC, distributor of Aleve®.

NDC 69517-109-24
HealthLife
Naproxen Sodium Caplets 220 mg
 Pain Reliever / Fever Reducer (NSAID)
 Strength to Last 12 Hours
 24 CAPLETS
 Compare to Aleve® active ingredient†

Distributed by: HealthLife of USA LLC
 Rahway, NJ 07065
 www.healthlifeofusa.com
 0 69517 10924 8
 Lot. No.:
 Exp. Date:

Drug Facts
Active ingredient (in each caplet)
 Naproxen sodium 220 mg (naproxen 220 mg) (NSAID)†. Pain reliever. Fever reducer. Nonsteroidal anti-inflammatory drug.
Uses Temporarily relieves minor aches and pains due to: minor pain of arthritis, backache, headache, muscular aches, menstrual cramps, toothache, common cold. Temporarily reduces fever.
Warnings Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.
Each bleeding warning: This product contains an NSAID. NSAIDs may increase the risk of bleeding. The chance is higher if you: are age 60 or older, take a blood thinning (anticoagulant) or steroid drug, take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), have 3 or more alcoholic drinks every day while using this product, take more or for a longer time than directed.
Do not use: If you have ever had an allergic reaction to any other pain reliever/fever reducer. Right before or after heart surgery.

Drug Facts (continued)
 doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center immediately.
Directions Do not take more than directed. The smallest effective dose should be used.
 Adults and children 12 years and older: Take 1 caplet every 8 to 12 hours while symptoms last. Do not take more than 2 caplets in any 8-12 hour period. Do not exceed 2 caplets in a 24-hour period.
 Children under 12 years: ask a doctor.
Other information Do not take with other NSAIDs (naproxen, ibuprofen, aspirin, etc.) or with alcohol. Store at 20°-25°C (68°-77°F). Avoid high humidity and excessive heat above 40°C (104°F).
KEEP OUT OF REACH OF CHILDREN. WARNING: UNDER CAP IS BROKEN OR MISSING
Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, FD&C Blue #2 lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, titanium dioxide.
Questions or comments? Call toll free 1-844-832-1138 Monday through Friday 9AM - 5PM EST or www.healthlifeofusa.com

Drug Facts (continued)
Ask a doctor before use: stomach bleeding warning applies to you; you have a history of stomach problems, such as heartburn you have high blood pressure, heart disease, liver cirrhosis, or kidney disease; you are taking a diuretic; you have problems or serious side effects from taking pain relievers or fever reducers; you have asthma.
Ask a doctor or pharmacist before use if you are: under a doctor's care for any serious condition; taking any other drug.
When using this product: take with food or milk if stomach upset occurs; the risk of heart attack or stroke may increase if you use more than directed or for longer than directed.
Stop use and ask a doctor if: you experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools; pain gets worse or lasts more than 3 days; fever gets worse or lasts more than 3 days; you have difficulty swallowing; it feels like the pill is stuck in your throat; redness or swelling is present in the painful area; any new symptoms appear.
If pregnant or breast-feeding: ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a

DESIGN DEPARTMENT, INC. - 716-873-5008

 DIE LINE DO NOT PRINT	 VARNISH AREA	Job prints 4 color process, please match	 CMYK
Date To Disk: 5/13/16 5/19/16	Job No.: 9382 9382	Plus 1 Spot Color	 PMS 7687

If Design Department, Inc. does not have an opportunity to view a "HARD COPY" or "PAPER" proof we will not be responsible for incorrect colors or printing errors in final printing. WE DO NOT CONSIDER ANY ELECTRONIC FILES INCLUDING PDF FILES AS A HARD COPY OR PAPER PROOF.


DRUG FACTS FORMAT WARNING
 Title - 9pt
 Headings - 8pt
 Body Copy - 5.5pt

We experienced problems formatting the Drug Facts information into the allotted space. Please note the above concessions to the minimum allowable text sizes.