



NDC 69517-105-24

Extra Strength Headache Relief

Acetaminophen 250 mg / Pain Reliever
Aspirin 250 mg (NSAID) / Pain Reliever
Caffeine 65 mg / Pain Reliever Aid



24 CAPLETS SEE NEW WARNINGS INFORMATION

Compare to Excedrin® Extra Strength active ingredients!

Distributed by: HealthLife of USA LLC
 Rahway, NJ 07065
www.healthlifeofusa.com



0 69517 10524 0

Lot. No.:
Exp. Date:

Drug Facts (continued)

Questions or comments?
 Call toll free 1-844-832-1138
 Monday through Friday 9AM - 5PM EST or
www.healthlifeofusa.com

RETAIN CARTON FOR COMPLETE DRUG FACTS

†This product is not manufactured or distributed by Novartis Consumer Health, Inc., owner of the registered trademark Excedrin® Extra Strength

Drug Facts (continued)

Take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen or others) **Do not use if you are taking a longer time than directed**

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, such as caffeine capsules, tea, coffee, energy drinks, sleepiness, and, occasionally, rapid heartbeat.

Do not use if you ever had an allergic reaction to acetaminophen, aspirin, or any other pain reliever/fever reducer (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have liver disease, stomach bleeding warning applies to you, such as heartburn, indigestion, or stomach problems, you have high blood pressure, heart disease, liver cirrhosis or kidney disease, you have asthma, you are taking a diuretic.

Ask a doctor or pharmacist before use if you are taking a prescription drug for diabetes, ulcers, or arthritis, taking any other drug or are under a doctor's care for any serious condition.

Stop use and ask a doctor if following signs of stomach bleeding occur: **black stools, vomit blood, or have stomach pain that does not get better.**

Drug Facts (continued)

Directions

Other information

Inactive ingredients

Drug Facts (continued)

Active ingredients Purpose (in each caplet)

Uses

Warnings

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older, have had stomach ulcers or bleeding problems, or are a blood thinning anticoagulant or steroid drug.

DESIGN DEPARTMENT, INC. - 716-873-5008

DIE LINE DO NOT PRINT	VARNISH AREA	pms 7482	pms 185	Process Yellow

Date To Disk: 5/13/16
5/19/16

Job No.: 9381
9381

Black

If Design Department, Inc. does not have an opportunity to view a "HARD COPY" or "PAPER" proof we will not be responsible for incorrect colors or printing errors in final printing. WE DO NOT CONSIDER ANY ELECTRONIC FILES INCLUDING PDF FILES AS A HARD COPY OR PAPER PROOF.

DRUG FACTS FORMAT WARNING

Title - 9pt
Headings - 8pt
Body Copy - 5pt

We experienced problems formatting the Drug Facts information into the allotted space. Please note the above concessions to the minimum allowable text sizes.